

CROSS-CULTURAL EXPERIENCE OF MATERNAL POSTNATAL DEPRESSION

¹Siti Roshaidai Mohd Arifin, ²Helen Cheyne, ³Margaret Maxwell

ABSTRACT---The prevalence of maternal postnatal depression (PND) varies from 0% to 60% globally. This wide variety brings up the issue of whether PND is a universal medical condition or whether it is an idea impacted by cultural and social translations, and the labelling of signs and symptoms. The objective of this review was to understand women's experience of PND in different countries. Studies reporting women's experiences of PND were searched through databases of CINAHL, PubMed, MEDLINE, PsycINFO and ASSIA databases using specific key words. Articles published between 2006 and 2016 were filtered for inclusion criteria. A total of 27 studies on maternal experience of PND conducted in ten different countries including America, Canada, South Africa, United Kingdom, Norway, Australia, New Zealand, Bangladesh, China, and Taiwan were reviewed. Findings indicated that while women recognized the emotional changes in themselves after their childbirth, they were unable to perceive these as burdensome symptoms, resulting in delayed diagnosis of PND. The issues of cultures and traditions were perceived by Asian women as one of the contributing factors to PND. HCPs were regarded by the women as having a lack of knowledge in supporting mental wellbeing among postnatal women. Therefore, it is crucial to educate both HCPs and communities to notice and react to women's depressed feelings. The management of maternal PND should acknowledge the social and cultural element as many women associated this with the development of PND.

Keywords---qualitative; mothers; perspectives; experiences; postnatal depression; maternal mental health.

I. Introduction

Maternal postnatal depression (PND) is a burdensome disease which most much of the time starts 2 to 3 weeks after birth and that may keep going for a year and may have considerable consequences for new moms, babies, accomplices and families.^(1,2) Without proper management, PND can cause adverse effects on mother, baby and partners, impacting familial relationships as well as the maternal ability to care for their baby.⁽³⁻⁵⁾

The wide variety of prevalence of PND (between 0% to 60%) in new mothers from various societies⁽⁶⁾ raises doubt about whether PND mirrors an all-inclusive therapeutic determination as characterized by the Diagnostic and Statistical Manual of Mental Disorders (DSM)- V, or an idea that is impacted by cultural and social elucidation and

¹Department of Professional Nursing Studies, International Islamic University Malaysia, Kuantan, Pahang, Malaysia.

^{2,3}Nursing, Midwifery and Allied Health Professionals (NMAHP) Research Unit, University of Stirling Scotland, United Kingdom.

Corresponding email: roshaidai@iium.edu.my

labelling of signs and symptoms.⁽⁷⁾ It was recommended that cultural and social elements should be incorporated in managing maternal PND,⁽⁸⁻¹¹⁾ therefore it is crucial to understand how women from different cultural backgrounds describe their experiences of PND and to what extent it affects their help seeking behaviours.

Whilst previous qualitative studies have explored the experience of women with depressive symptoms in various cultural backgrounds,⁽¹²⁻¹⁴⁾ limited studies have specifically examined women's experiences and perceptions of PND across different cultures. In their review, Dennis and Chung-Lee⁽¹⁵⁾ included 40 studies on maternal coping styles and ideal treatment in dealing with PND. Women in their review did not disclose their feelings due to several barriers. First, the women perceived that their family members and healthcare professionals (HCPs) were reluctant to respond to their emotional and practical needs. Second, the women had a lack of knowledge about PND, therefore were unable to recognize PND symptoms. Third, women were afraid of being labelled as mentally ill and of losing their right to care for their newborn.

Dennis et al.⁽¹⁶⁾ reviewed 51 studies on traditional postpartum practices in over 20 countries. They had found that there were commonalities and difference in practices across cultures. Regardless of their cultural backgrounds, women in their review reported that they were supported by their family members during the early postnatal period with restrictions on their freedom to perform their normal household tasks. In many cultures, women described restrictions on their behaviour, for example some food was prohibited and some permitted and they were prescribed certain hygiene practices (such as washing genitalia with soap and water, avoiding sexual intercourse).⁽¹⁶⁾ Despite the similarities, there were variations in terms of duration of postnatal practices, participants in the postnatal practices, and the influence of supernatural or religious beliefs.

Although issues of barriers to help-seeking and maternal treatment preferences, traditional postpartum practices and rituals were discussed in the previous review articles^(15,16) there are no cross-cultural reviews available on the women's description of the symptoms of PND, and their interpretation of the perceived causes of those symptoms. The aims of this review were to systematically examine whether there are cross cultural differences in maternal PND experience and in their coping styles, and to examine positioning of sociocultural factors as explanations for PND.

II. Methods and methodology

A qualitative synthesis of women's experience of PND was conducted. Qualitative synthesis allows researchers to employ systematic methods of scientific inquiry in order to aggregate or summarize the previous qualitative data.⁽¹⁷⁾ The outcome of this process was the generation of new themes. It is through this process that makes qualitative synthesis used in this review differs from literature reviews.

Papers included in this review based on the following criteria: using qualitative data collection and analysis; conducted among women with PND and published in English from year 2006 to 2016. Two previous reviews on PND relied on data reported in 2005 and July 2006.^(15,16) Only qualitative papers are considered in this review as the aim is to capture the experience of PND as expressed by the women, not based on the women's interpretation of the

specific set of depressive symptoms found from self-reported questionnaires to detect PND. The search strategy is shown in Figure 1. The Critical Appraisal Skills Program (CASP)⁽¹⁸⁾ was used not only to assess the quality but also to review methods and methodology the included papers. To guide the writing of the result of this review, the Enhancing Transparency in Reporting the Synthesis of Qualitative Research (ENTREQ) statement was adopted.

Papers selected for this review were analyzed using thematic analysis. Based on this type of analysis, repeating concepts or ideas from the selected papers were recognized and analyzed transparently.^(19, 20) The selected papers were exported to Nvivo for the purpose of data management and analysis. The interpretation and conclusion of this review were derived from three systematic steps. First, findings of the selected papers were read line-by-line and relevant ideas will be coded along with the reading. Second, the ‘descriptive themes’ were developed based on the coding in the first step. The coding was compared across the papers. Third, ‘descriptive themes’ were analyzed and the new construct (analytical themes) were developed.




Figure 1: Search strategy

III. Findings

A total of 27 selected studies came from ten different countries (America, Canada, South Africa, UK, Norway, Australia, New Zealand, and Bangladesh, China, and Taiwan). Almost all papers meeting most of the CASP criteria, indicating high quality of the papers. Results of this review were classified under three themes as indicated below.

Symptoms

The experience of PND was described by the women as a sudden change in their normal self, sadness, and oversensitive. They connected these experiences with emotional disturbance, tearfulness, so much pressure, feeling guilty, anxiety, and self-blaming.^(21–29) Women, especially those in the two studies from South Africa uncovered that they had short-tempered, feeling irritated, disturbing thoughts and self-destructive ideation. In one study, this irritability was converted into forceful conduct, for example, hitting kids and notwithstanding cutting their husband.⁽²⁴⁾ Women in Canada, South Africa, Taiwan, and Bangladesh reported difficulty in sleeping, being absent minded, weight reduction, absence of hunger, stomach torments, chest torments, migraine, tipsiness, low or hypertension, jaundice, urinary issues, gastric issues, and experiencing difficulty strolling after childbirth.^(22,24,26,27,30)

Women's attribution of the symptoms

The women' attribution of PND was essentially connected with their social circumstances which were accounted for by practically most of the studies. While women across cultures recognized biophysical factors as adding to PND, the essentialness of cultures and tradition were especially talked about in three Asian studies.^(22,30,31) Women linked their experience of PND with an absence of help from their informal community, money related issues, multiple roles, and problematic relationships which led them to feelings of isolation, fear, disappointment, powerlessness, and sentiments of being disregarded.^(21–24,26,29,30,32–39)

Financial constraints were discussed by many women from American, South African, Indian, Welsh, Chinese, Bangladeshi and immigrant women in the USA and Canada as one of the cause for their PND^(22–24,26,31,32,34,36,40) Various studies had found that women with PND experienced trouble in adapting to their different duties as another mother and put remarkable weight on them.^(21,25,29,31–35,38,40) The sense of inability to fulfil expected roles became significant when there was a mismatch between maternal desire and the truth of turning into a mother.^(21,26,29,31,33–36,39,42,43)

Problematic relationships were highlighted across different studies.^(22,24,26,30,32,33,37,42) While Australian women revealed that their partner was not understanding and unsympathetic towards the manner in which they were feeling and this led to disappointment, Asian migrant ladies in Canada depicted conflicts with their relative who they frequently portrayed as uncalled for or unsupportive, coinciding with PND.^(26,33)

Most of the women in this review linked their PND experience to unpleasant occasions during the postnatal period such as having breastfeeding problems, multiple roles, and insufficient rest and sleep.^(21–26,30–33,35,36,38,39) While women in western countries regarded their motherhood as 'less or no reward', 'being unable to escape', 'not worth',

and 'no-win' situation,^(23,33,35,36) Asian women described it as 'collapsing', 'cursing' 'feeling drained' and 'had no physical strength'.^(22,30,31)

Women's accounts indicated that their experience of PND were related to traumatic labour experiences, articulating the feeling that they 'couldn't really recover', 'stressful', and 'frightened during labour and delivery'.^(21-23,32,33) Women who had high expectations of breastfeeding appeared to describe that their high desire to continue breastfeeding increased their distress and gave them the sense of disappointment whenever they had breastfeeding problems.^(21,31,41)

In certain communities, traditional postnatal practices were reported as women from PND, yet it could likewise add to a negative effect in another culture. For instance, South African women perceived such practices as quiet, alleviating, helping with relief from discomfort, and advancing profound sleep.⁽²³⁾ Similarly, in Canada, a study of immigrant women suggested that the absence of the practices had decreased the women's supportive network which had exposed them to PND.^(26,40) However, many Asian women reported that their traditional rituals during postnatal period had contributed to their PND experience.^(30,31) For instance, the Chinese women portrayed the practice of remaining inside the house for a month during postnatal period as 'being in jail' and losing of authority over themselves.^(30,31)

The issues of gender preference of the newborn and the presence of an 'external power' were highlighted by Bangladeshi, Chinese and Indian women. They reported that they were blamed by their husband and in laws for not having the option to bring forth a male child.^(21,25,30) Some Asian women connected their PND experience with the presence of an external power and they described the power as 'evil eye', 'evil fortune', and 'satanic whisper'.^(22,28-30)

Help seeking behaviours

It appeared that women included within this review have limited understanding of PND. It is due to this lack of knowledge that made them unable to identify the symptoms, therefore embracing self-improvement measures to alleviate the symptoms.^(21,25-27,29,33,34,36-38,42,44,45) Those women who had a certain level of knowledge on depressive symptoms attempted to conceal their feelings because of not wanting being labelled as a weak or bad mother or being viewed 'differently' by their society, and to avoid the consequences of losing their when the referral to social services becomes necessary.^(23,25-27,29,33-37,40,42-45) A lack of knowledge of PND among family and HCPs were also reported by the women. They were described by the women as not being supportive in getting professional help and had normalized their depressive symptoms.^(21, 24-27,29,34-38,40,42-45) Nevertheless, some women chose to seek help when they were encouraged by their HCPs, spouse, relatives and friends to do so.^(25,27,45) Women portrayed the supportive HCPs as 'good listener', 'goddesses', and 'understanding and unrushed'.^(21,37,38,44)

Women used various strategies to improve their depressive conditions such as having social support, practicing self-help, and receiving professional support.^(26,37,38) Practicing self-help and receiving social support were discussed

more by Asian women, and they did not describe professional help at all.^(22,29,30,38) Spiritual beliefs were perceived by the immigrant women in Canada and South Asian women in England as beneficial in coping with the depressive symptoms, whereas women in Europe were bound to discuss perceiving their very own needs and individual alteration, for example, keeping occupied and escaping the house each day.^(21,29,39-41) Women in other western countries (Australia, New Zealand, UK, and US) appreciated the professional help such as psychologically informed sessions, listening visits, home visits, and telephone-based services in promoting their mental well-being.^(25,26,28, 33, 34, 36,37,41,44,45)

IV. Discussions

This review indicated that while social circumstances were equally discussed by all group of women as one of the main causes for PND, the issues of traditional postnatal practices were discussed more by Asian women. Similar to previous studies, this review recommends that non-performance of postnatal traditions and rituals has been inconsistently associated with PND and that performance of some postnatal traditions is related to PND.⁽⁴⁶⁻⁴⁹⁾ The trend towards modernization, migration and globalization may reduce the capability of a woman to follow tradition postnatal practices or to feel stressed when they are asked to follow the practices that they no longer believe.⁽¹⁶⁾

Many women included in this review were reported as having a lack of understanding of depressive symptoms, resulting in delayed diagnosis of PND. While they recognized the emotional changes in themselves after their childbirth, they were frequently unable to perceive these as burdensome symptoms. Similarly, Dennis and Chung-Lee⁽¹⁵⁾ suggested that the inadequate knowledge on PND had resulted in inability to recognize the symptoms of PND, therefore was considered as a major barrier for seeking professional help. Furthermore, women also regarded their HCPs as having a lack of understanding in supporting mental wellbeing among postnatal mothers. Therefore, it is crucial to educate HCPs and communities to acknowledge and react to maternal emotional changes after childbirth.⁽⁵⁰⁾

This finding suggests that supportive social networks are for many women across cultures.^(25,27,28,34,45) Within Asian communities, self-help was identified as the main coping strategies for PND. Similarly, Dennis and Chung-Lee⁽¹⁵⁾ summarized that women preferred to have ‘talking therapies’ rather than receive medication. It was suggested that the therapy should acknowledge the social and cultural element⁽¹¹⁾ as many women associated this with the development of PND.

There were two limitations of this review. First, only studies published in the previous 10 years were included in this review. Second, there were wide variations in methods used across the included studies such as PND qualitative design, assessment time frame for PND and focus of topic guide, creating challenges for comparing findings across studies. Despite these limitations, this review addresses the literature on the women’s experience of PND across cultures and their help seeking styles, collectively.

V. Conclusion

This review suggested that women across cultures have limited understanding of postnatal mental wellbeing, therefore commonly were not able to acknowledge their emotional disturbance after childbirth as PND. While women across cultures shared some similarities of their PND experiences, there were some culturally specific experiences that were not common in other parts of the world. For instance, Asian women associated their PND experience with the issue of cultures and traditions during postnatal period. This possibly due to the impact of an interpretive lens that makes PND being interpreted differently from one woman to another.⁽⁵¹⁾ This review recommends that the women's perceptions of their PND experience should be acknowledged within the clinical practice. For example, in offering the therapy for a woman with PND, HCPs could use the knowledge about the women's attribution of the symptoms to improve their adherence with the treatment plan.⁽⁵²⁾

Source of Funding: The authors would like to thank the International Islamic University Malaysia for funding this research through 'Research Initiative Grant Scheme' (RIGS16-142-0306).

Ethical Clearance: Not Applicable

Conflict of Interest: Nil

References

- [1] World Health Organisation. Maternal mental health and child development in low- and middle-income countries. 2008 [Accessed 19 December 2017]. Available from: http://www.who.int/mental_health/prevention/suicide/mmh_jan08_meeting_report.pdf
- [2] American Psychological Association (APA). Postpartum depression: What is postpartum depression and anxiety? [Internet]. 2014 [Accessed 2017 Dec 19]. Available from: <http://www.apa.org/pi/women/programs/depression/postpartum.aspx>
- [3] Brett K, Barfield W, Williams C. Prevalence of self-reported postpartum depressive symptom – 17 states, 2004–2005. 2008 Apr [Accessed 2017 Dec 19];57(14):361–66. Available from: <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5714a1.htm>
- [4] McGarry J, Kim H, Sheng X, Egger M, Baksh L. Postpartum depression and help-seeking behavior. *J Midwifery Womens Health*. 2009 Jan-Feb;54(1):50–6. Available from: <https://doi.org/10.1016/j.jmwh.2008.07.003>
- [5] Pitanupong J, Liabsuetrakul T, Vittayanont A. Validation of the Thai Edinburgh postnatal depression scale for screening postpartum depression. *Psychiatry Res*. 2007 Jan;149(1–3):253–259. Available from: <http://dx.doi.org/10.1016/j.psychres.2005.12.011>
- [6] Halbreich U, Karkun S. Cross-cultural and social diversity of prevalence of PPD and depressive symptoms. *J Affect Disord*. 2006 Apr;91(2–3):97–111. Available from: <http://doi.org/10.1016/j.jad.2005.12.051>
- [7] American Psychiatric Association (APA). Diagnostic and statistical manual of mental disorders (DSM) [Internet]. 2013 [Accessed 2017 Dec 19]. Available from: <http://psychgradaran.narod.ru/lib/clinical/DSM5.pdf>
- [8] Davy GM, Ehiobuche C. Health care and culture clash: The case of Caribbean immigrants in the US. *International Journal of Arts & Sciences*. 2013;6(2):501–14. Available from: <http://universitypublications.net/ijas/0602/pdf/H3V784.pdf>
- [9] Mamisachvili L, Ardiles P, Mancewicz G, Thompson S, Rabin K, Ross LE. Culture and postpartum mood problems: Similarities and differences in the experiences of first- and second-generation Canadian

- women. J Transcult Nurs. 2013 Jan;24(2):162–70. Available from: <https://doi.org/10.1177/1043659612472197>
- [10] Niemi ME, Falkenberg T, Nguyen MTT, Nguyen MTN, Patel V, Fazelid E. The social contexts of depression during motherhood: A study of explanatory models in Vietnam. *J Affect Disord*. 2010 Jul;124(1–2):29–37. Available from: <http://dx.doi.org/10.1016/j.jad.2009.09.017>
- [11] Rahman A. Challenges and opportunities in developing a psychological intervention for perinatal depression in rural Pakistan--a multi-method study. *Arch Womens Ment Health*. 2007 Oct;10(5):211–19. Available from: <http://dx.doi.org/10.1007/s00737-007-0193-9>
- [12] Bowen A, Duncan V, Peacock S, Bowen R, Schwartz L, Campbell D, et al. Mood and anxiety problems in perinatal Indigenous women in Australia, New Zealand, Canada, and the United States: A critical review of the literature. *Transcult Psychiatry*. 2014 Feb;51(1):93–111. Available from: <http://dx.doi.org/10.1177/1363461513501712>
- [13] Jackson F. 2013. Investigating women’s experiences of postpartum depression in Barbados. 141st APHA Annual Meeting, 2013 Nov 2–6, Boston, MA. Washington, DC: American Public Health Association. Available from: <https://apha.confex.com/apha/141am/webprogramadap/Paper283729.html>
- [14] Williams P. Mothers' descriptions of recovery from postpartum depression. *MCN Am J Matern Child Nurs*. 2013 Sep/Oct;38(5):276–81. Available from: <https://doi.org/10.1097/NMC.0b013e3182993fbf>
- [15] Dennis CL, Chung-Lee L. Postpartum depression help-seeking barriers and maternal treatment preferences: A qualitative systematic review. *Birth*. 2006 Dec;33(4):323–31. Available from: <http://doi.org/10.1111/j.1523-536X.2006.00130.x>
- [16] Dennis CL, Fung K, Grigoriadis S, Robinson GE, Romans S, Ross L. Traditional postpartum practices and rituals: A qualitative systematic review. *Women's Health*. 2007 Jul;3(4):487–502. Available from: <http://doi.org/10.2217/17455057.3.4.487>
- [17] Noblit G, Hare R. *Meta-ethnography: Synthesising qualitative studies*. 11th ed. London: Sage Publications; 1988.
- [18] Public Health Resource Unit. *The Critical Skills Appraisal Programme: making sense of evidence*. 2006. Public Health Resource Unit, England. Available: <http://www.casp-uk.net/>
- [19] Khan N, Bower P, Rogers, A. Guided self-help in primary care mental health: Meta-synthesis of qualitative studies of patient experience. *Br J Psychiatry*. 2007 Sep;191(3):206–11. Available from: <https://doi.org/10.1192/bjp.bp.106.032011>
- [20] Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol*. 2008 Jul;8:45. Available from: <https://doi.org/10.1186/1471-2288-8-45>
- [21] Coates R, Ayers S, de Visser R. Women's experiences of postnatal distress: A qualitative study. *BMC Pregnancy Childbirth*. 2014 Oct;14:359. Available from: <https://doi.org/10.1186/1471-2393-14-359>
- [22] Edhborg M, Nasreen HE, Kabir ZN. "I can't stop worrying about everything" – Experiences of rural Bangladeshi women during the first postpartum months. *Int J Qual Stud Health Well-being*. 2015 Jan;10(1):26226. Available from: <http://doi.org/10.3402/qhw.v10.26226>
- [23] Kathree T, Petersen I. South African Indian women screened for postpartum depression: A multiple case study of postpartum experiences. *S Afr J Psychol*. 2012 Mar;42(1):37–50. Available from: <https://doi.org/10.1177/008124631204200105>
- [24] Kathree T, Selohilwe OM, Bhana A, Petersen I. Perceptions of postnatal depression and health care needs in a South African sample: The “mental” in maternal health care. *BMC Womens Health*. 2014 Nov;14:140. Available from: <https://doi.org/10.1186/s12905-014-0140-7>
- [25] McCarthy M, McMahon C. Acceptance and experience of treatment for postnatal depression in a community mental health setting. *Health Care Women Int*. 2008 Jun;29(6):618–37. Available from: <https://doi.org/10.1080/07399330802089172>
- [26] Morrow M, Smith JE, Lai Y, Jaswal S. Shifting landscapes: immigrant women and postpartum depression. *Health Care Women Int*. 2008 Jun;29(6):593–617. Available from: <https://doi.org/10.1080/07399330802089156>
- [27] Sword W, Busser D, Ganann R, McMillan T, Swinton M. Women's care-seeking experiences after referral for postpartum depression. *Qual Health Res*. 2008 Sep;18(9):1161–73. Available from: <https://doi.org/10.1177/1049732308321736>
- [28] Williamson V, McCutcheon H. The experiences of Australian women diagnosed with postnatal depression who received health professional intervention. *SingapNurs J*. 2007;34:39–44.

- [29] Wittkowski A, Zumla A, Glendenning S, Fox JRE. The experience of postnatal depression in South Asian mothers living in Great Britain: A qualitative study. *J Reprod Infant Psych*. 2012 Jan;29(5):480–92. Available from: <https://doi.org/10.1080/02646838.2011.639014>
- [30] Chen CH, Wang SY, Chung UL, Tseng YF, Chou FH. Being reborn: The recovery process of postpartum depression in Taiwanese women. *J AdvNurs*. 2006 May;54(4):450–6. Available from: <http://dx.doi.org/10.1111/j.1365-2648.2006.03843.x>
- [31] Gao L, Chan SW, You L, Li X. Experiences of postpartum depression among first-time mothers in mainland China. *J AdvNurs*. 2010 Feb;66(2):303–12. Available from: <http://doi.org/10.1111/j.1365-2648.2009.05169.x>
- [32] Abrams LS, Curran L. “And you're telling me not to stress?” A grounded theory study of postpartum depression symptoms among low-income mothers. *Psychol Women Q*. 2009 Sep;33(3):351–62. Available from: <http://dx.doi.org/10.1111/j.1471-6402.2009.01506.x>
- [33] Buultjens M, Liamputtong P. When giving life starts to take the life out of you: Women's experiences of depression after childbirth. *Midwifery*. 2007 Mar;23(1):77–91. Available from: <http://dx.doi.org/10.1016/j.midw.2006.04.002>
- [34] Callister LC, Beckstrand RL, Corbett C. Postpartum depression and help-seeking behaviours in immigrant Hispanic women. *J ObstetGynecol Neonatal Nurs*. 2011 Jul–Aug;40(4): 440–9. Available from: <http://dx.doi.org/10.1111/j.1552-6909.2011.01254.x>
- [35] Dennis TR, Moloney MF. Surviving postpartum depression and choosing to be a mother. *South Online J Nur Res*. 2009[Accessed 2017 December 19];9(4). Available from: http://www.resourcenter.net/images/snrs/files/sojnr_articles2/Vol09Num04Art06.html
- [36] Hanley J, Long B. A study of Welsh mothers' experiences of postnatal depression. *Midwifery*. 2006 Jun;22(2):147–57. Available from: <http://doi.org/10.1016/j.midw.2005.08.004>
- [37] O'Mahony JM, Donnelly TT, Raffin BS, Este D. Cultural background and socioeconomic influence of immigrant and refugee women coping with postpartum depression. *J Immigr Minor Health*. 2013 Apr;15(2):300–14. Available from: <http://dx.doi.org/10.1007/s10903-012-9663-x>
- [38] Shafiei T, Small R, McLachlan H. Immigrant Afghan women's emotional well-being after birth and use of health services in Melbourne, Australia. *Midwifery*. 2015 Jul;31(7):671–7. Available from: <https://doi.org/10.1016/j.midw.2015.03.011>
- [39] Vik K, Hafting M. “Smile through it!” Keeping up the facade while suffering from postnatal depressive symptoms and feelings of loss: Findings of a qualitative study. *Psych*. 2012 Sep;3(9A):810–17. Available from: <https://doi.org/10.4236/psych.2012.329123>
- [40] O'Mahony JM, Donnelly TT, Raffin BS, Este D. Barriers and facilitators of social supports for immigrant and refugee women coping with postpartum depression. *AdvNurs Sci*. 2012 Jul/Sep;35(3):E42–56. Available from: <http://dx.doi.org/10.1097/ANS.0b013e3182626137>
- [41] Homewood E, Tweed A, Cree M, Crossley J. Becoming occluded: The transition to motherhood of women with postnatal depression. *Qual Res Psychol*. 2006 Nov;6(4):313–29. Available from: <https://doi.org/10.1080/14780880802473860>
- [42] Bilszta J, Ericksen J, Buist A, Milgrom J. Women's experience of postnatal depression – Beliefs and attitudes as barriers to care. *Aust J AdvNurs*. 2010 Mar/May;27(3):44–54. Available from: <https://search.informit.com.au/documentSummary;dn=909929185761432;res=IELHEA>
- [43] Chew-Graham C, Sharp D, Chamberlain E, Folkes L, Turner KM. Disclosure of symptoms of postnatal depression, the perspectives of health professionals and women: A qualitative study. *BMC Fam Pract*. 2009 Jan;10:7. Available from: <http://doi.org/10.1186/1471-2296-10-7>
- [44] Shakespeare J, Blake F, Garcia J. How do women with postnatal depression experience listening visits in primary care? A qualitative interview study. *J Reprod Infant Psych*. 2006 Jan;24(2):149–62. Available from: <https://doi.org/10.1080/02646830600643866>
- [45] Slade P, Morrell CJ, Rigby A, Ricci K, Spittlehouse J, Brugha TS. Postnatal women's experiences of management of depressive symptoms: a qualitative study. *Br J Gen Pract*. 2010 Nov;60(580):e440–448. Available from: <https://doi.org/10.3399/bjgp10X532611>
- [46] Chan SW, Levy V, Chung TK, Lee D. A qualitative study of the experiences of a group of Hong Kong Chinese women diagnosed with postnatal depression. *J AdvNurs*. 2002 Sep;39(6):571–9. Available from: <http://dx.doi.org/10.1046/j.1365-2648.2002.02326.x>
- [47] Fisher JRW, Morrow MM, Nhu Ngoc NT, Hoang Anh LT. Prevalence, nature, severity and correlates of postpartum depressive symptoms in Vietnam. *BJOG*. 2004 Dec;111(12):1353–60. Available from: <http://doi.org/10.1111/j.1471-0528.2004.00394.x>

- [48] Lee DT, Yip AS, Leung TY, Chung TK. Ethnoepidemiology of postnatal depression. Prospective multivariate study of sociocultural risk factors in a Chinese population in Hong Kong. *Br J Psychiatry*. 2004 Jan;184:34–40. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/14702225>
- [49] Rahman A, Iqbal Z, Harrington R. Life events, social support and depression in childbirth: perspectives from a rural community in the developing world. *Psychol Med*. 2003 Nov;33(7):1161–7. Available from: <https://doi.org/10.1017/S0033291703008286>
- [50] Knudson-Martin C, Silverstein R. Suffering in silence: A qualitative meta-data-analysis of postpartum depression. *J Marital Fam Ther*. 2009 Apr;35(2):145–58. Available from: <https://doi.org/10.1111/j.1752-0606.2009.00112.x>
- [51] Walsh D, Evans K. Critical realism: An important theoretical perspective for midwifery research. *Midwifery*. 2014 Jan;30(1):e1–e6. Available from: <https://doi.org/10.1016/j.midw.2013.09.002>
- [52] Jorm AF. Mental health literacy: Public knowledge and beliefs about mental disorders. *Br J Psychiatry*. 2000 Nov;177(5):396–401. Available from: <https://doi.org/10.1192/bjp.177.5.396>
- [53] Samsudeen, A., Sulphey, M.M. Does performance management of medical practitioners stand apart?(2018) *International Journal of Pharmaceutical Research*, 10 (2), pp. 101-106. <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85043774593&partnerID=40&md5=4cead8e5a3b57c125044af857bf04a60>