

# POSTPARTUM HAEMORRHAGE IN MATERNAL MOTHERS AT ANUTAPURA PUBLIC HOSPITAL IN PALU, 2017

<sup>1</sup>Rosmala Nur, <sup>2</sup>Hasra Sartika Sarina, <sup>3</sup>Nurhaya S Patui, <sup>4</sup>Sitti Radhiah, <sup>5</sup>Novi Inriyanny Suwendro, <sup>6</sup>Ariani, <sup>7</sup>Syahrir A Pasinringi

**ABSTRACT**--Postpartum haemorrhage is one of the factors causing maternal mortality in Indonesia. In Indonesia, at least 128.000 women suffer from haemorrhage causing them to death. Data from the Anutapura Public Hospital in Palu showed that the rate of postpartum haemorrhage within 2015-2017 was fluctuated and still caused maternal mortality. This study aimed to determine the risk factors of the incidence of postpartum haemorrhage in maternal mothers at Anutapura Public Hospital, Palu. The type of study was observational with a case-control approach. The subject of the study was mothers having a postpartum haemorrhage. The total sample of this study was 50 people as the case group and 100 people as the control group by adjusting the types of labour. Sampling was done using a total sampling technique. The data were analyzed using an odds ratio test with  $\alpha = 5\%$ . This study showed that anemia (OR=2.874 and CI=1.421-5.812), parity (OR=3.995 and CI=1.952-8.174), age (OR=2.874 and CI=1.421-5.812) were the risk factors of postpartum haemorrhage in women. It can be concluded that the anemia, parity, and age are the risk factors of postpartum haemorrhage in women. The importance of increasing ANC (antenatal care) visits is to control maternal and fetal health and provide information through a medical consultation about the importance of planning a good pregnancy.

**Keywords**--Postpartum Haemorrhage, Anemia, Parity, Age.

## I. INTRODUCTION

The success of maternal health attempts can be seen from the indicator of Maternal Mortality Ratio (MMR). MMR is the total maternal mortality during pregnancy, labour, and puerperium caused by pregnancy, labour, and puerperium or the management, but it is not because of the other causes such as accident or fall per 100,000 live births. Then, maternal mortality can be changed into the ratio of maternal mortality and it is expressed per 100,000 live births by dividing the mortality rate by the general fertility rate. By using this method, it can obtain a maternal mortality ratio per 100,000 births (Survey. 2017).

The maternal mortality rate is one of the vital indicators for global Millenium Development Goals (MDGs) that had ended in 2015, which was continued by the Sustainable Development Goals (SDGs) up to 2030. According

---

<sup>1</sup> Public Health Department, Faculty of Public Health, Universitas Tadulako, Palu, Indonesia, nurrosmla09@gmail.com

<sup>2</sup> Public Health Department, Faculty of Public Health, Universitas Tadulako, Palu, Indonesia

<sup>3</sup> Public Health Department, Faculty of Public Health, Universitas Tadulako, Palu, Indonesia

<sup>4</sup> Public Health Department, Faculty of Public Health, Universitas Tadulako, Palu, Indonesia

<sup>5</sup> Public Health Department, Faculty of Public Health, Universitas Tadulako, Palu, Indonesia

<sup>6</sup> Nutrition Department, Faculty of Public Health, Universitas Tadulako, Palu, Indonesia

<sup>7</sup> Faculty of Public Health, Hasanuddin University, Makassar, Indonesia

to the interpretation of the goals and the global SDGs for the indicators of a healthy and prosperous life, in 2030, the maternal mortality rate should be decreased by up to less than 70 per 100,000 live births (*Sastrawinata* 2015).

According to the WHO report in 2014, the Maternal Mortality Ratio in the world was 289 per 100,000 live births, 179 per 100,000 live births for North Africa, and 16 per 100,000 live births for Southeast Asia. The Maternal Mortality Ratio in Asian countries was as follows 214 per 100,000 live births for Indonesia, 170 per 100,000 live births for the Philippines, 160 per 100,000 live births for Vietnam, 44 per 100,000 live births for Thailand, 60 per 100,000 live births for Brunei, and 39 per 100,000 live births for Malaysia. Alternatively, WHO estimated that in 2015, the total maternal mortality reached around 303 per 100,000 live births globally. Maternal Mortality Ratio (MMR) in some developing countries reached 239 per 100,000 live births that were 20 times greater than that of the developed countries. The developing countries contributed around 90% or 302 per 100,000 live births from the estimated maternal mortality ratio that happened in 2015 (*Perempuan et al.* 2019).

The profile of Health Agency in Central Sulawesi in 2018 showed the primary four causes for maternal deaths including haemorrhage of 30.3%, pre-eclampsia of 27.1%, infection of 7.3%, and other factors that caused a direct maternal death of 35.3%, such as cancer, kidney disease, heart disease, or other diseases suffered by the mothers (*Sulawesi.* 2018).

Based on the data collected from the Health Agency of Palu City in the annual report within 2015 - 2017, it could be found a case of maternal mortality caused by haemorrhage, whereby in 2015, it was collected 22 cases of maternal mortality with 326/100,000 live births, 11 cases of maternal mortality with 158/100,000 live births in 2016, and 11 cases of maternal mortality with 156/100,000 live births in 2017. Besides the data of mortality rate, there was data about the factors causing maternal mortality from 2015 to 2017; the factors causing maternal mortality were haemorrhage of 63.63% followed by pre-eclampsia of 22.72%, infection of 4.5%, and other factors of 9.09%. Moreover, the maternal mortality rate based on the place of death was shown as follows; 66.30% at Regional Public Hospital (*RSUD*), which Anutapura Public Hospital in Palu was the place with the highest maternal mortality rate, namely 5 deaths out of 11 deaths in 2017 (*Palu* 2015).

The data from Anutapura Public Hospital in Palu showed that the incidence rate of postpartum haemorrhage in 2015 was 36 cases from 3,412 birth mothers, 58 cases from 3,243 birth mothers in 2016, while in 2017, the rate was 50 cases from 2,804 birth mothers. Even though there was a decline, postpartum haemorrhage still becomes the key factor causing a high rate of maternal mortality (*Department of Health* 2016) (*Dinas kesehatan provinsi sulawesi Tengah* 2019).

Most maternal mortality cases happen 24 hours after giving birth and postpartum haemorrhage is one of the direct causes of maternal mortality. Postpartum haemorrhage is a leading unexpected cause of maternal mortality in the world. The haemorrhage case in Indonesia is estimated at around 14 million. There is, at least, 128,000 women experiencing haemorrhage until it causes deaths annually. Postpartum haemorrhage is the most frequent factor causing maternal mortality. It is the post-delivery bleeding happened within the first 24 hours following childbirth (*Maswime and Buchmann* 2016).

Postpartum haemorrhage is excessive bleeding in the genital tract after the birth of a baby or abortion and until 6 weeks after that. The primary postpartum haemorrhage refers to bleeding within 24 hours following birth, while the secondary postpartum haemorrhage is the bleeding occurred after 24 hours (and in 6 weeks).<sup>7</sup> Many risk factors cause postpartum haemorrhage, such as having a pregnancy complication grade 3 in the medical history previously,

previous cesarean section, multiple pregnancy, primigravida, high parity, anemia, surgical labour, prolonged obstructed labour, precipitate parturition, placenta previa, abruptio placentae (Yuliati *et al.* 2018). This study aimed to know the risk factors of the incidence of postpartum haemorrhage in birth mothers at Anutapura Public Hospital, Palu, in 2017.

## II. METHOD

The type of this study was a case-control study with an analytical survey as the approach to know the risk factors of the dependent variable and independent variable. The dependent variable in this study was postpartum haemorrhage, while the independent variable was the factors might risk resulting in the incidence of postpartum haemorrhage, namely Anemia, Parity, and age.

The type of data in this study was secondary data. The data in this study were collected from the patients' medical records having complete data in 2017. The data analysis was done using univariate and bivariate analyses by presenting the results in the form of tables.

## III. RESULTS

The result of bivariate analysis for the risk factors of the incidence of postpartum haemorrhage in birth mothers at Anutapura Public Hospital, Palu, in 2017 is as below.

**Table 1 The Risk of Anemia and the Incidence of Postpartum Haemorrhage**

Anemia	Postpartum Haemorrhage				Total	OR (CI 95%)
	Case		Control			
	F	%	F	%		
High Risk	27	54.0	138	13.0	35	7.856
Low Risk	23	46.0	7	87.0	115	(3.510 –
<b>Total</b>	<b>50</b>	<b>100.0</b>	<b>100</b>	<b>100.0</b>	<b>150</b>	<b>17.582)</b>

*Source: Primary Data, 2017*

Table 1 shows the respondents with postpartum haemorrhage are those with a high risk of anemia; the total is 54.0% that is higher than the total patients with a low risk of anemia of 46.0%. Meanwhile, the respondents with no postpartum haemorrhage are mostly non-anemic of 87.0% that is higher than the total respondents who are anemic of 46.0%.

Based on the statistical test, it was collected an OR value of 7.8 at 95% CI of 3.510 – 17.582. It indicated that the risk of postpartum haemorrhage in anemic birth mothers was 7.8 times greater than that of non-anemic birth mothers. It is because the OR value was more than 1 and the lower limit and the upper limit were more than 1, so anemia was the significant risk factor in postpartum haemorrhage.

**Tabel 2 The Risk of Parity and the Incidence of Postpartum Haemorrhage**

Parity	Postpartum Haemorrhage		Total	OR (CI 95%)
	Case	Control		

	<b>F</b>	<b>%</b>	<b>F</b>	<b>%</b>		
High Risk	31	62.0	297	29.0	60	3.995
Low Risk	19	38.0	1	71.0	90	(1.952– 8.174)
<b>Total</b>	<b>50</b>	<b>100.0</b>	<b>100</b>	<b>100.0</b>	<b>150</b>	

*Source: Primary Data, 2017*

Table 2 shows that the total respondents with postpartum haemorrhage having parity of more than 3 are 62.0% and the total of those having parity of around 2-3 is 38.0%. Meanwhile, the total respondents with no Postpartum haemorrhage having parity of around 2-3 is 71.0% and the total respondents having parity of more than 3 is 29.0%.

Based on the statistical test, it was obtained an OR value of 3.9 at 95% CI of 1.952 – 8.174, indicating that the risk of postpartum haemorrhage in birth mothers with parity of more than 3 is 3.9 times greater than the birth mothers having parity of around 2 to 3. It is because the OR value was more than 1 and the lower limit and the upper limit were more than 1, so parity was the significant risk factor in postpartum haemorrhage.

**Table 3 The Risk of Age and the Incidence of Postpartum Haemorrhage**

<b>Age</b>	<b>Postpartum Haemorrhage</b>				<b>Total</b>	<b>OR (CI 95%)</b>
	<b>Case</b>		<b>Control</b>			
	<b>F</b>	<b>%</b>	<b>F</b>	<b>%</b>		
High Risk	27	54.0	297	29.0	59	2.874
Low Risk	23	46.0	1	71.0	94	(1.421 – 5.812)
<b>Total</b>	<b>50</b>	<b>100.0</b>	<b>100</b>	<b>100.0</b>	<b>150</b>	

*Source: Primary Data, 2017*

Table 3 shows the total respondents with Postpartum haemorrhage in mothers at the age of below 20 years old and over 35 years old is higher (54.0%) than that of the mothers at the age of around 20 to 35 years old (46.0%). Meanwhile, the total respondents with no postpartum haemorrhage in the mothers at the age of below 20 years old and over 35 years old (71.0%) is higher than that of the mothers at the age of around 20 to 35 years old (29.0%).

Based on the statistical test, it was obtained an OR value of 2.8 at 95% CI of 1.421 – 5.812, indicating that the risk of postpartum haemorrhage in birth mothers at the age of below 20 years old and over 35 years old is 2.8 times greater than that of the birth mothers at the age of around 20 to 35 years old. It is because the OR value was more than 1 and the lower limit and upper limit were more than 1, so Age was the significant risk factor of postpartum haemorrhage.

## IV. Discussion

### *Anemia and the Risk of Postpartum haemorrhage*

Anemia can weaken the mothers' immunity and increase the frequency of complications while pregnant and labour. Anemia also causes an increased risk of postpartum haemorrhage.<sup>8</sup> Getting tired easily in anemic patients is caused by the energy metabolism by the muscles that cannot move perfectly due to lack of oxygen. Based on the result of the analysis of the risk of anemia in this study, it was obtained an Odds Ratio (OR) with 95% Confidence Interval (CI) of 7.8, indicating that the anemic birth mothers have the risk of postpartum haemorrhage of 7.8 times greater than the non-anemic birth mothers (Karlsson *et al.* 2015).

These research findings are in line with the study conducted by (*Sentilhes et al.* 2015) that the anemic birth mothers have a risk of postpartum haemorrhage 4.8 times greater, indicating that there is a relationship between anemia and the incidence of postpartum haemorrhage in birth mothers.

During pregnancy, more iron is needed to produce red blood cells because the mothers should fulfill the fetal needs and themselves; also, during the labour process, the mothers need hemoglobin intake to provide energy, so the uterine muscle can have a good contraction (*Dupont et al.* 2009).

### ***Parity and the Risk of Postpartum haemorrhage***

The mothers with high parity will affect the condition of the mothers' uterus because the more frequent the mothers to give birth, the reproductive function decreases, the uterine muscle will be too tight, and it cannot have a normal contraction; so, it allows a high possibility of having postpartum haemorrhage (*Maryani et al.* 2016).

Based on the result of analysis for the risk of parity in this study, it was obtained an Odds Ratio (OR) with 95% Confidence Interval (CI) of 3.9, indicating that the birth mothers with parity of more than 3 had a risk of postpartum haemorrhage 3.9 times greater than that of the birth mothers with parity of around 2-3. In this study, it was recorded that from 50 postpartum haemorrhage cases, 31 of them were the mothers with parity of more than 3 (high risk), while the total anemic mother with hemoglobin levels of less than 11 g/dl (high risk) was 27 and the total mother at the age of below 20 years old and over 35 years old (high risk) was 27 people.

The research findings showed that the total respondent experiencing postpartum haemorrhage with parity of more than 3 (62.0%) was higher than those with parity of around 2-3 (38.0%), while the total respondent with no Postpartum haemorrhage having parity of around 2-3 (71.0%) was higher than those with parity of more than 3 (29.0%).

The result of this study is in line with (*Cortet et al.* 2012) and (*Janighorban et al.* 2018) stating that primipara and multipara women lose their blood as much as in postpartum haemorrhage; in this case, multipara women lose their blood more than nulliparous women (nullipara). Parity has a role in causing the incidence of postpartum haemorrhage because, in each pregnancy and labour, there is a change in the uterine muscle fiber that can decrease the uterus ability to have a contraction. Consequently, it is difficult to press the opened blood vessels after the placenta detaches. The risk of having the incidence will increase after the third labour or more causing postpartum haemorrhage.

### ***Age and the Risk of Postpartum haemorrhage***

The best age for women to get pregnant with low risk is around 20 to 35 years old because they are in a healthy reproductive phase. Maternal mortality in pregnant mothers and birth mothers at the age of below 20 years old and over 35 years old will significantly increase because they are exposed to the complication both medical and obstetric; it can harm the mothers' lives; it is the reason why age becomes the cause of postpartum haemorrhage (*Yuliati et al.* 2018) (*Natarajan et al.* 2016). In this study, it was recorded that from 50 postpartum haemorrhage cases, 27 of them were the mothers at the age of below 20 years old and over 35 years old (high risk). Also, the total anemic mother with hemoglobin levels of less than 11g/dl was 27 people (high risk) and the total mother with parity of more than 3 (high risk) was 31 people. The result of this study showed that the respondents experiencing postpartum haemorrhage were mostly mothers at the age of below 20 years old and over 35 years old (54.0%). The

total was higher than that of the mothers at the age of around 20 to 35 years old (46.0%). The respondents with no postpartum haemorrhage were mostly mothers at the age of around 20 to 35 years old (71.0%). It was higher than mothers at the age of below 20 years old and over 35 years old (29.0%). Based on the result of the analysis for risk of age in this study, it was collected an Odds Ratio (OR) with 95% Confidence Interval (CI) of 2.8, indicating that birth mothers at the age of below 20 years old and over 35 years old had a risk of postpartum haemorrhage 2.8 times greater than the birth mothers at the age of around 20 to 35 years old.

The result of this study is in line with the study conducted by (Collins *et al.* 2017)(Quantin *et al.* 2017) that birth mothers at the age of below 20 years old and over 35 years old have a higher risk of postpartum haemorrhage. It is also in line with (Bai *et al.* 2010) that age is strongly connected with the readiness of the mothers in reproduction. Women at the age of below 20 years old are still at the growth and development stages, so the pregnancy condition will make them should share with the fetus they are expecting to fulfill the nutrition needs. Contrary, mothers at the age of over 35 years old start to show the effect of aging, such as frequently having a disease like hypertension and diabetes mellitus that can inhibit fetal food entering through a placenta. Besides, postpartum haemorrhage causing maternal mortality in pregnant women that gives birth at the age of below 20 years old is 2.5 times greater than postpartum haemorrhage happened at the age of around 20 years old to 35 years old. Postpartum haemorrhage increases again after the age of around 30-35 years old (Mamar *et al.* 2020).

## V. CONCLUSION

Based on the study above, it can be concluded that anemia, parity, and mothers' age are the risk factor of the incidence of postpartum haemorrhage in birth mothers. Anutapura Public Hospital of Palu City is expected to make the result of this study as a reference in designing the preventive program in birth mothers having a risk factor to experience postpartum haemorrhage and the mitigation program for the incidence of postpartum haemorrhage, so it can decrease the complication caused by the incidence.

## REFERENCES

1. Bai, J., Wong, F. W. S., Bauman, A., and Mohsin, M. (2010). "Parity and pregnancy outcomes," *American Journal of Obstetrics and Gynecology*, 186(2), 274–278. DOI: 10.1067/mob.2010.119639
2. Collins, P. W., Bruynseels, D., Mallaiah, S., Dick, J., Elton, C., Weeks, A., Sanders, J., Aawar, N., Townson, J., Hood, K., and Hall, J. (2017). "Viscoelastometry guided fresh frozen plasma infusion for postpartum haemorrhage : OBS2 , an observational study," *British Journal of Anaesthesia*, The Author(s), 119(3), 422–434. DOI: 10.1093/bja/aex245
3. Cortet, M., Dupont, C., Colin, C., and Rudigoz, R. (2012). "Association between fibrinogen level and severity of postpartum haemorrhage : secondary analysis of a prospective trial," *British Journal of Anaesthesia*, *British Journal of Anaesthesia*. Published by Elsevier Ltd., 108(6), 984–989. DOI: 10.1093/bja/aes096
4. Department of Health. (2016). *Profile of Health in Central Sulawesi Province*, Department of Health of Central Sulawesi Province, Palu.
5. Dinas kesehatan provinsi sulawesi Tengah. (2019). "Profil kesehatan Provinsi Sulawesi tengah Tahun

- 2019,” 1–222.
6. Dupont, C., Touzet, S., Colin, C., Rabilloud, M., Clement, H. J., Lansac, J., Colle, M. H. B., and Rudigoz, R. C. (2009). “Incidence and management of postpartum haemorrhage following the dissemination of guidelines in a network of 16 maternity units in France,” *International Journal of Obstetric Anesthesia*, Elsevier Ltd, 18(4), 320–327. DOI: 10.1016/j.ijoa.2009.02.017
  7. Janighorban, M., Heidari, Z., and Dadkhah, A. (2018). “Women ’ s Needs on Bed Rest during High-risk pregnancy and Postpartum Period : A Qualitative Study,” *Midwifery and Reproductive Health*, 2(1), 1–9. DOI: 10.22038/jmrh.2018.28162.1304
  8. Karlsson, O., Jeppsson, A., Thornemo, M., Lafrenz, H., Rådström, M., and Hellgren, M. (2015). “Fibrinogen plasma concentration before delivery is not associated with postpartum haemorrhage : a prospective observational study,” 115(March), 99–104. DOI: 10.1093/bja/aev039
  9. Mamar, S., Puspasari, I., Demak, K., B, E. Y., Rusydi, M., and R, E. T. (2020). “Mother ’ s Behavior in Pregnancy-Puerperal Treatments and Reproductive Health Disorders.”
  10. Maryani, S., Respati, S. H., and Astirin, O. P. (2016). “Association Between Pregnant Woman Class and Pregnancy Complication in Tegal District, Central Java,” *Journal of Maternal and Child Health*, 01(04), 214–219. DOI: 10.26911/thejmch.2016.01.04.02
  11. Maswime, S., and Buchmann, E. (2016). “Causes and avoidable factors in maternal death due to cesarean-related hemorrhage in South Africa,” *International Journal of Gynecology and Obstetrics*, International Federation of Gynecology and Obstetrics, 134(3), 320–323. DOI: 10.1016/j.ijgo.2016.03.013
  12. Natarajan, A., Alaska, A., Nelson, B. D., Ahn, R., Oguttu, M., Dulo, L., Eckardt, M. J., and Burke, T. F. (2016). “International Journal of Gynecology and Obstetrics Provider experiences with improvised uterine balloon tamponade for the management of uncontrolled postpartum hemorrhage in Kenya,” *International Journal of Gynecology and Obstetrics*, International Federation of Gynecology and Obstetrics, 135(2), 210–213. DOI: 10.1016/j.ijgo.2016.05.006
  13. Palu, D. K. (2015). *Peserta KB Aktif menurut Metode Kontrasepsi Cara Modern Kota Palu pada Bulan Januari sampai Desember 2015*, Kota Palu.
  14. Perempuan, K. P., Anak, D. A. N. P., and Statistik, B. P. (2019). “Profil anak indonesia 2019.”
  15. Quantin, C., Sagot, P., Marocchini, M., and Laufe, J. (2017). “ScienceDirect Postpartum hemorrhage with transfusion : Trends , near misses , risk factors and management at the scale of a perinatal network ´ chelle d ’ un re,” 46, 455–460. DOI: 10.1016/j.jogoh.2017.03.011
  16. Sastrawinata, U. S. (2015). “Optimalisasi Persalinan Non-institusional Untuk Menurunkan Angka Kematian Ibu Era Millenium Development Goals Optimalization Non-institutional Delivery in Reducing Maternal Mortality Ratios To Achieve Millenium Development Goals,” 41(8), 212–219. DOI: <http://dx.doi.org/10.15395/mkb.v41n4.256>
  17. Sentilhes, L., Lasocki, S., Deruelle, P., Dreyfus, M., Perrotin, F., and Goffinet, F. (2015). “Tranexamic acid for the prevention and treatment of postpartum haemorrhage,” *British Journal of Anaesthesia*, The Author(s), 114(4), 576–587. DOI: 10.1093/bja/aeu448
  18. Sulawesi., H. D. of C. (2018). *The Profile of Health of Central Sulawesi Province, Yearly Report of Health Department of Central Sulawesi*, Palu.
  19. Survey., I. D. and H. (2017). *Jakarta Indonesia. ORC Macro*. Calverton, Maryland, USA.
  20. Yuliati, A., Soejoenoes, A., Suwondo, A., Anies, A., and Kartasurya, M. I. (2018). “Beberapa Faktor

Kejadian Perdarahan Postpartum Ibu Bersalin yang Dirawat Di Rumah Sakit,” Jurnal Epidemiologi  
Kesehatan Komunitas, 3(1), 7. DOI: 10.14710/jekk.v3i1.3101