

AWARENESS OF INTERFERON THERAPY AMONG DENTAL STUDENTS

Nithyanandham Masilamani¹, Dhanraj Ganapathy²

Abstract

Interferon is perhaps the term given to antiviral material formed by many vertebrate cells in reaction to viral infections. Research has established a number of mechanisms involved, in therapeutic benefits of interferon together with antiviral, innate immunity-enhancing and cytostatic functions. This survey was performed for assessing the awareness about interferon therapy amongst dental students. A cross sectional survey was performed with a self-administered questionnaire with 10 queries circulated among 100 dental students. The questionnaire assessed the awareness about Interferon therapy, their medicinal uses, anticancer activity, mechanism of action and side effects. The responses were recorded and analysed. 91% of the respondents were not aware of medical uses of Interferon therapy. 93% were not aware of anticancer activity of Interferon therapy. 93% were not aware of the mechanism of action of Interferon therapy. 96% were not aware of side effects of Interferon therapy. The awareness about the use of interferon therapy is very less among dental students. Increased awareness programs and sensitization programs should be conducted to improve the awareness levels.

Keywords: Awareness, interferon therapy, dental students

Introduction

Interferon is perhaps the term given to antiviral material formed by many vertebrate cells in reaction to viral infections. In 1957, Isaacs and Lindeman identified a component conferring the property of viral interruption (Isaacs & Lindenmann, 1987). The word interferon (IFN) has been created to describe the whole new substance. During the next 40 years, IFNs have been tested as treatment methods in a wide range of benign and malignant conditions. Research has established a number of mechanisms involved, in therapeutic benefits of interferon together with antiviral, innate immunity-enhancing and cytostatic functions

¹ Tutor, Department of Prosthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Chennai, India.

² Professor & Head of Department, Department of Prosthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Chennai, India, Email: dhanrajmaganapathy@yahoo.co.in

IFNs are an effective therapy in oncology for a variety of firm tumors and hematological cancers , t hat also involve skin cancer, renal cell carcinoma, AIDS-related Kaposi tumor , follicular lymphoma, acute myeloid leukemia, and chronic myelogenous leukemia (CML). IFN effects have been linked with negative side effects that have an effect on the standard of life of the patient and the capacity of the practitioner to efficiently manage the patient (Isaacs & Lindenmann, 1987).

IFNs were classified as alpha (α), beta (β), and gamma (Δ), referring to the former designations of leukocyte, fibroblast and immune IFN (Nomenclature, & Mouse., 1980.). Until lately, IFNs have been divided into two main subtypes due to its capability to connect to different receptor forms (Aguet et al . , 1984; Merlin et al . , 1985). Type I IFNs all connect to IFN type I receptors, which also include IFN- α , IFN- β , IFN- α , which IFN- α . IFN- Δ is the only type II IFN and connects to different type II receptor. (Walter et al., 1998).

The knowledge of IFN biochemistry, interactions, evidence for use, side-effect characteristics as well as the monitoring of IFN toxicity will contribute to the effective usages of these substances in the treatment of cancer patients. Interferon therapy has the possibility for use as a treatment modality for oral cancer. Hence,this survey was performed for assessing the awareness about interferon therapy amongst dental students

Materials and method

A cross sectional survey was performed with a self-administered questionnaire with 10 queries circulated among 100 dental students. The questionnaire assessed the awareness about Interferon therapy,their medicinal uses,anticancer activity,mechanism of action and side effects.The responses were recorded and analysed

Results

91% of the respondents were not aware of medical uses of Interferon therapy (Fig 1) .93% were not aware of anticancer activity of Interferon therapy (Fig 2) .93 % were not aware of the mechanism of action of Interferon therapy (Fig 3) .96% were not aware of side effects of Interferon therapy(Fig 4) .

Fig.1:Awareness of medical uses of Interferon therapy




Fig.2: Awareness of anticancer activity of Interferon therapy




Fig.3: Awareness of mechanism of action of Interferon therapy





Fig.4: Awareness of side effects of Interferon therapy



Discussion

The IFNs have a wide range of movement and are engaged with complex associations. They show antiviral movement, sway cell digestion and separation, and have antitumor action. The antitumor impacts have all the earmarks of being because of a mix of direct antiproliferative, just as circuitous insusceptible intervened impacts.

Interferons were extremely convincing in treating Hairy Cell Leukemia. The first evidence of IFN response to hairy cell leukemia were reported in 1984 (Quesada et al., 1984; Walter et al., 1998). Seven patients were treated with 3×10^6 U of part-filtered leukocyte IFN-5-007 i.m. Day after day (qd). Three patients had a total decrease (CR), four had an incomplete decrease (PR) and decreases were sustained for 6 to 10 months. In this way, 30 patients with leukemia were managed with IFN-5-0072a, including seven patients who had not been treated. Nine (30 per cent) CR and 17 (56 per cent) PR were confirmed by bone marrow biopsies.. Both hematological reports of patients either enhanced or consolidated (Aguet et al., 1984). A multi - center phase II analysis of 64 patients allocated in 1986 confirmed the efficacy of drugs for hairy cell leukemia (Golomb et al., 1986)

IFN-gm IFN-gm was used as a steady stage CML therapy. In 1987, Kurzrock et al. recorded that 6 out of 30 patients with unrelenting stage CML compensated with rIFN-gm showed CHR and oscillating degrees of cytogenetic improvement (Kurzrock et al . , 1987). Actually results considered that the use of IFN-5-007 and IFN- Δ mixtures has been ignored to demonstrate progress over IFN-5-007 alone and IFN- Δ is not recommended for diagnosis for CML. (KIoKe et al., 2009)

IFN has also been commonly used for the treatment of renal cell carcinoma. Regardless of the fact that there really is no reasonable position for its use in the adjuvant environment, extensive research has been done to define its role in the management of metastatic disease.(Pyrhönen et al., 1999)In 1980, Bart et al. gave an account of the restriction of B16 melanoma in vivo and in vitro via murine IFN (Bart et al., 1971). IFN monotherapy in patients with metastatic melanoma induces responses in nearly 15 % of cases.(Evans & Anthony Evans, 1992)

IFN is being used in the management of HIV-related KS from 1981 (Real et al., 1986). Initial medication regimens used 20×10^6 U / d doses associated with high response rates, and furthermore substantial levels of harmfulness.(Ward, 1986) In subsequent investigations (Dittmer & Krown, 2010), the probability of response to therapy was attributed to CD4 controls. Patients with CD4 sizes greater than $400 / \text{mm}^3$ responded, although none of those with CD4 tests of less than $150 / \text{mm}^3$ responded.

Adverse effect of interferons could be divided into strong and persistent indications. In an acute environment, patients experience fevers, chills and rigors, often in the 3-and 6-hour range after receiving IFN. Patients can also experience migraine headaches, myalgias, and pain. Resistance to these side effects can be triggered by the drawn out and continuous structure of IFN. Nevertheless , treatment intervals as short as a few days will lead to a regeneration of rigors and chills. Transaminitis and neutropenia may occur during an initial period of that not longer treatment duration and can be limited by modifying the IFN component. Both are decided by an infinite supply of medication. If transaminitis is not treated carefully, it can lead to deadly hepatic injury (Kirkwood et al., 1996). The constant side effects encountered by patients on IFN include fatigue (70%-100% of patients), anorexia (40%-70%) and neuropsychiatric manifestations (30%). Such adverse effects tend, in all instances, to be portion-related, and absolute, to be exacerbated after some period.

IFN is a promising however not entirely comprehended anticancer specialist. Clinical preliminary studies have provided different indications for IFNs across both hematological and potent tumor sectors, despite the fact that the appropriateness of IFNs is confidential in most of the operator's uses.The symptom patterns of the IFNs are substantial and, at large doses, the decision of the IFN treatment must be considered against the near conviction of the decrease in personal gratification of the patient.(Kirkwood et al., 1996; Sawyer et al., 1990)This study inferred the awareness about interferon therapy among dental students is poor.

Conclusion

The awareness about the use of interferon therapy is very less among dental students.Increased awareness programs and sensitization programs should be conducted to improve the awareness levels.Additional emphasis should be given in the curriculum and syllabi to incorporate the therapeutic benefits and upcoming research perspective of interferon therapy among dental students.This will enable and empower the students with the knowledge about all the versatile applications of interferon therapy in various therapeutic applications.

Funding Support :

The authors declare that they have no funding support for this study.

Conflict of Interest :

The authors declare that they have no conflict of interest.

References

1. Aguet, M., Gröbke, M., & Dreiding, P. (1984). Various human interferon a subclasses cross-react with common receptors: Their binding affinities correlate with their specific biological activities. In *Virology* (Vol. 132, Issue 1, pp. 211–216). [https://doi.org/10.1016/0042-6822\(84\)90105-3](https://doi.org/10.1016/0042-6822(84)90105-3)
2. Bart, R. S., Kopf, A. W., & Silagi, S. (1971). Inhibition of the Growth of Murine Malignant Melanoma by Polyinosinic-Polycytidylic Acid. In *Journal of Investigative Dermatology* (Vol. 56, Issue 1, pp. 33–38). <https://doi.org/10.1111/1523-1747.ep12291892>
3. Dittmer, D. P., & Krown, S. E. (2010). *Molecular Basis for Therapy of AIDS-Defining Cancers*. Springer Science & Business Media.
4. Evans, E. A., & Anthony Evans, E. (1992). Biologic therapy of cancer. Editors V T DeVita (Jr), S Hellman and S A Rosenberg Pub: J B Lippincott company, Philadelphia, USA, 1991, pp 816 price: US \$125.00 ISBN 0-397-51027-6. In *Journal of Labelled Compounds and Radiopharmaceuticals* (Vol. 31, Issue 1, pp. 79–80). <https://doi.org/10.1002/jlcr.2580310111>
5. Golomb, H. M., Jacobs, A., Fefer, A., Ozer, H., Thompson, J., Portlock, C., Ratain, M., Golde, D., Vardiman, J., & Burke, J. S. (1986). Alpha-2 interferon therapy of hairy-cell leukemia: a multicenter study of 64 patients. In *Journal of Clinical Oncology* (Vol. 4, Issue 6, pp. 900–905). <https://doi.org/10.1200/jco.1986.4.6.900>
6. Isaacs, A., & Lindenmann, J. (1987). Virus interference. I. The interferon. In *Journal of Interferon Research* (Vol. 7, Issue 5, pp. 429–438). <https://doi.org/10.1089/jir.1987.7.429>
7. KIoke, O., Wandl, U., Opalka, B., Moritz, T., Nagel-Hiemke, M., Franz, T., Becher, R., Hirche, H., Seeber, S., & Niederle, N. (2009). A prospective randomized comparison of single-agent interferon (IFN)-alpha with the combination of IFN-alpha and low-dose IFN-gamma in chronic myelogenous leukaemia. In *European Journal of Haematology* (Vol. 48, Issue 2, pp. 93–98). <https://doi.org/10.1111/j.1600-0609.1992.tb00572.x>
8. Kirkwood, J. M., Strawderman, M. H., Ernstoff, M. S., Smith, T. J., Borden, E. C., & Blum, R. H. (1996). Interferon alfa-2b adjuvant therapy of high-risk resected cutaneous melanoma: the Eastern Cooperative Oncology Group Trial EST 1684. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 14(1), 7–17.
9. [Kurzrock, R., Talpaz, M., Kantarjian, H., Walters, R., Saks, S., Trujillo, J. M., & Gutterman, J. U. \(1987\). Therapy of chronic myelogenous leukemia with recombinant interferon- gamma. In *Blood* \(Vol. 70, Issue 4, pp. 943–947\). <https://doi.org/10.1182/blood.v70.4.943.bloodjournal704943>](#)

10. Merlin, G., Falcoff, E., & Aguet, M. (1985). 125I-labelled Human Interferons Alpha, Beta and Gamma: Comparative Receptor-binding Data. In *Journal of General Virology* (Vol. 66, Issue 5, pp. 1149–1152). <https://doi.org/10.1099/0022-1317-66-5-1149>
11. Nomenclature, O., & Mouse, H. (1980). Interferon nomenclature. *Nature*, 286, 10.
12. Pyrhönen, S., Salminen, E., Ruutu, M., Lehtonen, T., Nurmi, M., Tammela, T., Juusela, H., Rintala, E., Hietanen, P., & Kellokumpu-Lehtinen, P.-L. (1999). Prospective Randomized Trial of Interferon Alfa-2a Plus Vinblastine Versus Vinblastine Alone in Patients With Advanced Renal Cell Cancer. In *Journal of Clinical Oncology* (Vol. 17, Issue 9, pp. 2859–2859). <https://doi.org/10.1200/jco.1999.17.9.2859>
13. Quesada, J. R., Reuben, J., Manning, J. T., Hersh, E. M., & Gutterman, J. U. (1984). Alpha Interferon for Induction of Remission in Hairy-Cell Leukemia. In *New England Journal of Medicine* (Vol. 310, Issue 1, pp. 15–18). <https://doi.org/10.1056/nejm198401053100104>
14. Real, F. X., Oettgen, H. F., & Krown, S. E. (1986). Kaposi's sarcoma and the acquired immunodeficiency syndrome: treatment with high and low doses of recombinant leukocyte A interferon. In *Journal of Clinical Oncology* (Vol. 4, Issue 4, pp. 544–551). <https://doi.org/10.1200/jco.1986.4.4.544>
15. Sawyer, L. A., Metcalf, J. A., Zoon, K. C., Boone, E. J., Kovacs, J. A., Clifford Lane, H., & Quinnan, G. V. (1990). Effects of interferon- α in patients with aids-associated Kaposi's sarcoma are related to blood interferon levels and dose. In *Cytokine* (Vol. 2, Issue 4, pp. 247–252). [https://doi.org/10.1016/1043-4666\(90\)90024-n](https://doi.org/10.1016/1043-4666(90)90024-n)
16. Walter, M. R., Bordens, R., Nagabhusan, T. L., Williams, B. R. G., Herberman, R. B., Dinarello, C. A., Borden, E. C., Trotta, P. P., Pestka, S., & Pfeffer, L. M. (1998). Review of Recent Developments in the Molecular Characterization of Recombinant Alfa Interferons on the 40th Anniversary of the Discovery of Interferon. In *Cancer Biotherapy and Radiopharmaceuticals* (Vol. 13, Issue 3, pp. 143–154). <https://doi.org/10.1089/cbr.1998.13.143>
17. Ward, T. T. (1986). *AIDS and Other Medical Problems in the Male Homosexual*.