

AN OVERVIEW ON ORAL MUCOCELE

Ghayathri Vijaya Kumaran¹, Nithya Jagannathan²

Abstract: *Mucoccele is a common lesion of the oral mucosa that results from an alteration of minor salivary glands due to a mucous accumulation. The most frequent localizations of these lesions are the lower lip mucosa. It also presents on other locations like tongue, buccal mucosa, soft palate, retromolar pad and lower labial mucosa. Two histological types exist, extravasation and retention. Mucocelles can appear at any site of the oral mucosa where minor salivary glands are present. Mucocelles can affect the general population but most commonly young patients between the age of 20-30 years. Clinically, they consist of soft bluish and transparent cystic swelling which occurs spontaneously.*

Keywords: *Mucoccele, minor salivary glands, extravasation, retention and buccal mucosa.*

1. INTRODUCTION:

The former is a swelling of connective tissue consisting of a collection of fluid called mucin. This occurs because of a ruptured salivary gland duct usually caused by local trauma (damage), in the case of mucus extravasation phenomenon, and an obstructed or ruptured salivary duct (Parotid) in the case of a mucus retention cyst. The mucoccele has a bluish translucent colour, and is more commonly found in children and young adults.[1] Although the term cyst is often used to refer to these lesions, mucocelles are not strictly speaking true cysts because there is no epithelial lining. Rather, it would be more accurate to classify mucocelles as polyps for an example, lump. There are two types of mucoccele which can appear as extravasation and retention. Extravasation mucoccele results from a broken salivary glands duct and the consequent spillage into the soft tissues around the gland.[2] Retention mucoccele appears due to decrease or absence of glandular secretion produced by a blockage of the salivary gland ducts. When located on the floor of the mouth and these lesions are called Ranulas. They are rarely found on the upper lip.[3] As their name suggests they are basically mucus lined cysts and they can also occur in the Paranasal sinuses most commonly the frontal sinuses, the frontoethmoidal region and also in the maxillary sinus.[4] Sphenoid sinus involvement is extremely rare. When the lumen of the vermiform appendix gets blocked due to any factor, again a mucoccele can form.

2. Discussion:

Oral mucocelles are asymptomatic lesions in most cases, but, in a few instances, when appearing as multiple and recurrent lesions they may cause severe pain. The lower lip is the most common site, although they can be found in any region where there are salivary glands, but rarely in the palate, retromolar space and the buccal mucosa.[5] Mucus is produced exclusively by the minor salivary glands and is also the most important substance secreted by the major

¹Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Chennai, India. E mail: ghayathri2795@gmail.com

²Senior Lecturer, Department of Oral Pathology, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Chennai, India.

Corresponding Author: Dhanraj Ganapathy

Professor and Head, Department of Prosthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Chennai, India, Email: dhanraj@saveetha.com

sublingual salivary gland. Mucocoeles can appear by an extravasation or retention mechanism. Extravasation mucocoele are caused by a leaking fluid from surrounding acini.

This type of mucocoele is commonly found on the minor salivary glands. Moreover, physical trauma can cause a leakage of salivary secretion into surrounding submucosal tissues.[6] Inflammation is obvious due to stagnant mucus resulting from extravasation.

3. Characteristics:

The size of oral mucocoeles vary from 1 mm to several centimetres and they usually are slightly transparent with a blue tinge.[7] On palpation, mucocoeles may appear fluctuant but can also be firm. Their duration lasts from days to years, and may have recurrent swelling with occasional rupturing of its contents. Mucocoele is the common salivary gland disorder and it is the second most common benign soft tissue tumour in the oral cavity .[8] They are characterized by accumulation of mucoïd material with rounded, well circumscribed transparent, bluish coloured lesions of variable size. It is a soft and fluctuant asymptomatic swelling with rapid onset which frequently resolves spontaneously.[9] Common in the lower lip but may occur in other locations also. The bluish discolouration is mainly due to the vascular congestion and cyanosis of the tissue above and the fluid accumulation below. It also depends on the size of the lesion and proximity to the surface and upper tissue elasticity. They are usually doom shaped swellings with intact epithelium.

4. Symptoms:

Mucocoeles often show up on the inside of your lower lips, your gums, the roof of your mouth, or under your tongue. Those on the floor of the mouth are called ranulas. [10]Mucocoele may often go unnoticed, however some individuals may report certain symptoms that may arise due to discomfort. [11]However in most cases, the condition is painless.

- A small bump most often under the tongue is observed. A bump may also be observed on other areas, along the floor of the mouth. The bump is often translucent or bluish in colour and often hard to touch.
- The swollen bump may interfere with the ability of the individual to swallow food. In case the bump is large enough it may also interfere with the individual's speech and include chewing food.

In some cases, the condition may be associated with excessive salivation in the mouth.

There are two phenomenon's which explain the etiology of oral mucocoele:

1. Mucus extravasation phenomenon explains oral mucocoele arising from the rupture in the salivary gland that resulted from trauma and thereby resulting in swelling of the connective tissue that composed of the machine that accumulated.
2. Mucus retention cyst explains the onset of oral mucocoele as a result of obstruction or rupture of the salivary gland.

5. Causes:

It is usually caused due to obstruction or rupture of a salivary gland duct in the mouth, which results in the swelling of a part of the gland.[12] There are few causes of oral mucocoele which is commonly known.

- Mouth injury may result in the damage to the duct, which in turn may result a mucocele
- Habitual biting of the lip and other mouth areas may also increase the risk of injury to the duct.
- Gum infection or disease of the oral cavity can result in infection of the duct, which in turn may obstruct the duct and form a mucocele.

The trauma in the salivary gland may be the result of the following:

- Mechanical injury during the process of mastication where the tissue of the lower lip gets caught between the teeth in the maxillary anterior and in the mandibular anterior.
- Habitual biting of the lip may accidentally bite the salivary gland resulting in trauma subsequently the onset of oral mucocele.
- Crush type injury can also result to oral mucocele
- Chronic irritation of the oral cavity such as from cigarette smoking and extreme heat can injure the salivary gland.
- Surgical procedures in the oral cavity may also cause trauma to the salivary gland.
- Congenital mucocele in newborn babies is believed to be the result of birth trauma affecting the oral cavity.

6. Treatment:

Conventional surgical removal is a most common method used to treat this lesion. Other treatment options include CO₂ laser ablation, cryosurgery, intralesional corticosteroid injection, micro marsupialization, marsupialization and electrocautery.[13] Extravasation and retention type of mucoceles clinically appears the same so that the treatment option for both the types are same. Moreover, small sized mucoceles are removed with marginal glandular tissues and in case of large lesions marsupialization needs to be taken. Lacrimal catheters are used to dilate the duct to remove the obstruction of retention type of mucoceles. [14] Other than that, if the fibrous wall of the mucocele is thick, then the removed tissue must be sent for histopathological examination to examine any salivary gland neoplasms. The micro marsupialization was considered as ideal and most commonly used treatment in case of pediatric patient because this technique was simple, rapid and prognosis is good.[15] The advantage in a CO₂ laser is that it minimizes the relapses and complications and allows rapid, simple mucocele ablation. It is also indicated for the patients who cannot tolerate long procedures.

Acknowledgement

The authors of this review acknowledge the institute, for their help towards collecting all the records and other data in relevance to the review article.

References:

1. Barankin B, Stedman TL, Metelitsa AI, Lin AN. Stedman's Illustrated Dictionary of Dermatology Eponyms. Lippincott Williams & Wilkins; 2005.
2. Bentley JM, Barankin B, Guenther LC. A Review of Common Pediatric Lip Lesions: Herpes Simplex/Recurrent Herpes Labialis, Impetigo, Mucoceles, and Hemangiomas. *Clinical Pediatrics* 2003;42:475–82. <https://doi.org/10.1177/000992280304200601>.
3. Har-El G, Balwally AN, Lucente FE. Sinus mucoceles: is marsupialization enough? *Otolaryngol Head Neck Surg* 1997;117:633–40.
4. Chi AC, Lambert PR 3rd, Richardson MS, Neville BW. Oral mucoceles: a clinicopathologic review of 1,824 cases, including unusual variants. *J Oral Maxillofac Surg* 2011;69:1086–93.

5. Tran TA, Parlette HL 3rd. Surgical pearl: removal of a large labial mucocele. *J Am Acad Dermatol* 1999;40:760–2.
6. [Jensen JL. Superficial Mucoceles of the Oral Mucosa. The American Journal of Dermatopathology 1990;12:88–92. https://doi.org/10.1097/00000372-199002000-00013.](#)
7. [Sugerman PB, Savage NW, Young WG. Mucocele of the anterior lingual salivary glands \(glands of Blandin and Nuhn\): Report of 5 cases. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology 2000;90:478–82. https://doi.org/10.1067/moe.2000.108805.](#)
8. [Ishii N, Maeyama Y, Nakama T, Hashimoto T. A case of solitary collagenoma localized on the upper lip mimicking mucocele. Clinical and Experimental Dermatology 2009;34:243–4. https://doi.org/10.1111/j.1365-2230.2008.02754.x.](#)
9. Silva A Jr, Nikitakis NG, Balciunas BA, Meiller TF. Superficial mucocele of the labial mucosa: a case report and review of the literature. *Gen Dent* 2004; 52:424–7.
10. Wood NK, Goaz PW. Differential Diagnosis of Oral and Maxillofacial Lesions. Mosby; 1997.
11. Yamasoba T, Tayama N, Syoji M, Fukuta M. Clinicostatistical study of lower lip mucoceles. *Head Neck* 1990;12:316–20.
12. Guimarães MS, Hebling J, Filho VAP, Santos LL, Vita TM, Costa CAS. Extravasation mucocele involving the ventral surface of the tongue (glands of Blandin-Nuhn). *Int J Paediatr Dent* 2006;16:435–9.
13. [Bermejo-Fenoll A, López-Jornet P. Instrument for Biopsy of Oral Lesions. Dermatologic Surgery 2006;32:1493–5. https://doi.org/10.1097/00042728-200612000-00011.](#)
14. [Ata-Ali J, Carrillo C, Bonet C, Balaguer J, Penarrocha M, Penarrocha M. Oral mucocele: review of the literature. Journal of Clinical and Experimental Dentistry 2010:e18–21. https://doi.org/10.4317/jced.2.e18.](#)
15. [Ma Y, Meng X, Su Y, Yan Z, Shao Q, Chen Y. The Bipedicle Advancement Flap on Skin Coverage After Mucous Cyst Excision:A Retrospective Study of 18 Cases n.d. https://doi.org/10.21203/rs.2.19837/v2.](#)